

# Explicit Bounds for the Burgess Bound for Character Sums

Enrique Treviño

Maine/Québec Number Theory Conference, 2009

# Short Character Sums

Let  $\chi$  be a non-principal character of modulus  $p$ .

$$S_{\chi}(N) = \sum_{M < n \leq N+M} \chi(n)$$

# Polya-Vinogradov

- Polya-Vinogradov  $S_\chi(N) \ll \sqrt{p} \log p$
- Constant made explicit and improved by people over time.
- Constant made explicit with a small constant and secondary term (Pomerance):

$$S_\chi(N) \leq \frac{1}{3 \log 3} \sqrt{p} \log p + 2\sqrt{p}$$

# Burgess

In the 60s, Burgess came out with the following:

## Theorem (D. Burgess)

Let  $\chi$  be a primitive character of conductor  $q > 1$ . Then

$$S_{\chi}(N) = \sum_{M < n \leq M+N} \chi(n) \ll N^{1-\frac{1}{r}} q^{\frac{r+1}{4r^2} + \epsilon}$$

for  $r = 2, 3$  and for any  $r \geq 1$  if  $q$  is cubefree, the implied constant depending only on  $\epsilon$  and  $r$ .

# Applications

- Improving upperbound for least quadratic non-residue (mod  $p$ )
- Calculating  $L(1, \chi)$

## Theorem (Iwaniec-Kowalski-Friedlander)

Let  $\chi$  be a Dirichlet character mod  $p$ . Then for  $r \geq 2$

$$|S_{\chi}(N)| \leq 30 \cdot N^{1-\frac{1}{r}} p^{\frac{r+1}{4r^2}} (\log p)^{\frac{1}{r}}.$$

## Improvement

## Theorem (ET)

Let  $\chi$  be a Dirichlet character mod  $p$ . Then for  $2 \leq r \leq 87$  and  $p \geq 10^7$ .

$$|S_\chi(N)| \leq 3 \cdot N^{1-\frac{1}{r}} p^{\frac{r+1}{4r^2}} (\log p)^{\frac{1}{r}}.$$

Note, the constant gets better for larger  $r$ , for example for  $r = 3, 4, 5, 6$  the constant is 2.376, 2.085, 1.909, 1.792 respectively.

## Proof

Idea 1: Shift, take average and use induction

$$S_{\chi}(N) = \sum_{M < n \leq M+N} \chi(n+ab) + \sum_{M < n \leq M+ab} \chi(n) - \sum_{M+N < n \leq M+N+ab} \chi(n)$$

$$1 \leq a \leq A, 1 \leq b \leq B.$$

Take average as  $a$  and  $b$  move around their options.



## Proof Cont.

$$V = \sum_{a,b} \sum_{M < n \leq M+N} \chi(n + ab)$$

Since  $\chi(n + ab) = \chi(a)\chi(\bar{a}n + b)$ , we have that

$$V = \sum_{x \pmod{p}} v(x) \left| \sum_{1 \leq b \leq B} \chi(x + b) \right|$$

where  $v(x)$  is the number of ways of writing  $x$  as  $\bar{a}n$  where  $a$  and  $n$  are in the proper ranges.

## Proof Cont.

## Idea 2: Holder's Inequality

- Let  $V_1 = \sum_{x \pmod{p}} v(x) = AN$
- Let  $V_2 = \sum_{x \pmod{p}} v^2(x)$
- Let  $W = \sum_{x \pmod{p}} \left| \sum_{1 \leq b \leq B} \chi(x+b) \right|^{2r}$ .

By Holder's Inequality we get

$$V \leq V_1^{1-\frac{1}{r}} V_2^{\frac{1}{2r}} W^{\frac{1}{2r}}$$

## Proof Cont.

## Lemma

For  $A \geq 40$  and  $A \leq \frac{N}{15}$ ,

$$V_2 = \sum_x \sum_{(\text{mod } p)} v^2(x) \leq 2AN \left( \frac{AN}{p} + \log(2A) \right)$$

$V_2$  is the number of quadruples  $(a_1, a_2, n_1, n_2)$  with  $1 \leq a_1, a_2 \leq A$  and  $M < n_1, n_2 \leq M + N$  such that  $a_1 n_2 \equiv a_2 n_1 \pmod{p}$ .

$$V_2 \leq AN + \sum_{a_1 < a_2} \left( \frac{(a_1 + a_2)N}{\gcd(a_1, a_2)p} + 1 \right) \left( \frac{\gcd(a_1, a_2)N}{\max\{a_1, a_2\}} + 1 \right)$$

## Ending Proof

- To bound  $W$  we use Weil's bound.
- Optimize the choices of  $A$  and  $B$ .
- Combine the lemmas with the induction hypothesis and figure out the constant.

# Quadratic Case (Booker)

## Theorem (Booker)

Let  $p > 10^{20}$  be a prime number  $\equiv 1 \pmod{4}$ ,  $r \in \{2, \dots, 15\}$  and  $0 < M, N \leq 2\sqrt{p}$ . Let  $\chi$  be a quadratic character  $\pmod{p}$ . Then

$$\left| \sum_{M \leq n < M+N} \chi(n) \right| \leq \alpha(r) p^{\frac{r+1}{4r^2}} (\log p + \beta(r))^{\frac{1}{2r}} N^{1-\frac{1}{r}}$$

where  $\alpha(r), \beta(r)$  are given by

$r$	$\alpha(r)$	$\beta(r)$	$r$	$\alpha(r)$	$\beta(r)$
2	1.8221	8.9077	9	1.4548	0.0085
3	1.8000	5.3948	10	1.4231	-0.4106
4	1.7263	3.6658	11	1.3958	-0.7848
5	1.6526	2.5405	12	1.3721	-1.1232
6	1.5892	1.7059	13	1.3512	-1.4323
7	1.5363	1.0405	14	1.3328	-1.7169
8	1.4921	0.4856	15	1.3164	-1.9808

## Improving Booker

For  $r \geq 3$  we can do the following:

$$c_r N^{1-\frac{1}{r}} p^{\frac{r+1}{4r^2}} \log(p)^{\frac{1}{2r}} < c_2 N^{\frac{1}{2}} p^{\frac{3}{16}} \log(p)^{\frac{1}{2}}$$

Then

$$N \leq \left( \frac{c_2}{c_r} \right)^{\frac{2r}{r-2}} p^{\frac{3r+2}{8r}} (\log(p))^{\frac{r-1}{r-2}}$$

Therefore we have  $N < \sqrt{p}$ . Hence the range Booker gets can be extended for  $r \geq 3$ .

# Improving the Log Factor in the General Case

The same trick gets us to improve my theorem to:

## Theorem (ET)

*Let  $\chi$  be a Dirichlet character mod  $p$ . Then for  $r \geq 3$ ,  $r \leq 74$  and  $p \geq 10^7$ .*

$$|S_{\chi}(N)| \leq 2.4 \cdot N^{1-\frac{1}{r}} p^{\frac{r+1}{4r^2}} (\log p)^{\frac{1}{2r}}.$$

# Acknowledgements

- My advisor Carl Pomerance for suggesting the problem and guidance.
- NSF, University of Maine for the financial support.
- Burgess, Iwaniec, Friedlander, Kowalski, Booker, Montgomery, Vaughan and Granville for their work on character sums.